

POST-OP INSTRUCTIONS FOR INFANT LIP & TONGUE-TIE RELEASE

Your goal is to have the frenum heal and re-form as far back as possible. You should do the stretches with the baby laying down on a bed or couch facing away from you like during the exam. You will receive a video link that demonstrates stretches. Please follow-up within 7-10 days. **Begin doing the stretches the DAY OF the procedure.** Gloves (preferred) or clean hands with nails trimmed should be used for stretches.

STRETCHES:

LIP-TIE RELEASE:

If the lip was revised, first put your fingers all the way in the fold of the lip and pull the lip up and out as high as possible, so you can see the white diamond and cover the nostrils. It may bleed slightly the first day or two, this is not a concern.

TONGUE-TIE RELEASE:

Option 1: Every 6 hours- Place one pointer finger underneath tongue and push back and down **slowly** as you count to 10. Repeat 2 more times.

Option 2: Every diaper change (every 2-3 hours)- Place one pointer finger underneath tongue and push back and down **slowly** as you count to 10. Repeat 1 time. It may bleed slightly the first day or two, this is not a concern.

- a. The main issue is to open the “diamond” all the way up on the lip and especially the tongue. If you notice it is becoming tight, then stretch a little more to open it back up as soon as possible.
- b. At other times, play in your child’s mouth a few times a day with clean fingers to avoid causing an oral aversion. Tickle the lips, the gums, or allow your child to suck your finger.
- c. Tummy-Time as much as possible. Visit www.TummyTimeMethod.com for helpful tips.

****WHAT TO EXPECT****

1. The released area will form a wet scab after the first day. It will appear white and soft. It may change color to yellow or even green. This is not infection, but is just a scab in the mouth. The white / yellow area will get smaller each day lengthwise, but HEALING IS STILL HAPPENING! So even though the white scab will heal you must continue stretching or the new frenum will not be as long as possible and the surgery may need to be repeated. If you have any concerns, please contact our office.
2. Your child’s lip will swell up slightly that evening or the next day. It is normal and will go down after a day or two. The area will be sore for a few days, at one week look much better, and at two weeks look much better and almost normal.
3. For pain control, you can give CHILDREN’S TYLENOL (160mg / 5mL) starting WHEN YOU GET HOME and for the next 2-3 days every 4-6 hours. Please give the recommended dose from your most recent pediatrician visit or based on the manufacturer’s recommended dosage for their age and weight. If your baby is refusing to nurse or seems to be in pain, please make sure the Tylenol dose is correct.
4. Follow up with a lactation consultant is critical if you are nursing. Bottle-feeding babies will benefit from visiting a feeding therapist. A bodyworker (chiropractor, CST, etc.) is also very helpful. You should expect one better feed a day (two better feeds the second day, etc.). Sometimes there’s an immediate difference in feeding, and sometimes it takes a few days. Skin to skin, warm baths, and soothing music can be very beneficial to calm the baby.

If you have any questions, please call us at 863-640-2150